Geometric classification of nilpotent Jordan algebras of dimension five

Iryna Kashuba[a,1,*], María Eugenia Martin[a,2]

[a]*Instituto de Matemática e Estatística, Universidade de São Paulo, R. do Matão 1010, 05508-090, São Paulo, Brazil.*

arXiv:1609.05594v1 [math.RA] 19 Sep 2016

Abstract

The variety $\mathcal{J}\mathrm{or}\mathrm{N}_5$ of five-dimensional nilpotent Jordan algebras structures over an algebraically closed field is investigated. We show that $\mathcal{J}\mathrm{or}\mathrm{N}_5$ is the union of five irreducible components, four of them correspond to the Zariski closure of the GL_5-orbits of four rigid algebras and the other one is the Zariski closure of an union of orbits of infinite family of algebras, none of them being rigid.

Keywords: Jordan algebra, deformation of algebras, rigidity
2010 MSC: 14J10, 17C55, 17C10.

1 Introduction

In this paper we study the geometry of nilpotent Jordan algebras over algebraically closed fields $\mathbf{k}$ of characteristic $\neq 2, 3$. Namely, let V be an n-dimensional $\mathbf{k}$-vector space, the bilinear maps $V \times V \to V$ form an n^3-dimensional algebraic variety $\mathrm{Hom}_{\mathbf{k}}(V \otimes V, V)$. Moreover, Jordan algebra structures satisfy polynomial equations (originating from both commutativity and Jordan identity) and therefore determine a Zariski-closed subset, which we will call the variety of Jordan algebras of dimension n. As any affine variety, $\mathcal{J}\mathrm{or}_n$ admits decomposition in a finite number of irreducible components. Furthermore, nilpotency conditions are also of polynomial type and thus the set of nilpotent Jordan algebras in dimension n, which will be denoted by $\mathcal{J}\mathrm{or}\mathrm{N}_n$, possess a structure of algebraic variety embedded in $\mathcal{J}\mathrm{or}_n$. The general linear group $\mathrm{G} = \mathrm{GL}(\mathrm{V})$ acts on both $\mathcal{J}\mathrm{or}_n$ and $\mathcal{J}\mathrm{or}\mathrm{N}_n$ via "change of basis", decomposing them into G-orbits which correspond to the classes of isomorphic Jordan algebras. We study the geometry of $\mathcal{J}\mathrm{or}\mathrm{N}(n)$: dimension, decomposition into irreducible components and their description. The crucial tool for it is the notion of deformation of algebras: we say that $\mathcal{J}_1$ is a deformation of $\mathcal{J}_2$ if the G-orbit of $\mathcal{J}_2$ belongs to Zariski closure of the G-orbit of $\mathcal{J}_1$. Algebra $\mathcal{J}$ whose orbit is an open subset in $\mathcal{J}\mathrm{or}_n$ is of particular interest, since it does not admit non-trivial deformations and its closure rises to irreducible component. We will call such algebras rigid.

Analogously to $\mathcal{J}\mathrm{or}\mathrm{N}_n$ one defines the varieties of nilpotent Lie and associative algebras. We will denote them $\mathrm{Lie}\mathrm{N}_n$ and $\mathrm{Assoc}\mathrm{N}_n$, respectively. The geometry of these two varieties is rather different.

*Corresponding author
Email addresses: kashuba@ime.usp.br (Iryna Kashuba), eugenia@ime.usp.br (María Eugenia Martin)
[1]The author was supported by CNPq (309742/2013-7).
[2]The author was supported by the CAPES scholarship for postdoc program of Mathematics, IME-USP.

In 1968, F. Flanigan posed a question whether every irreducible component is the closure of a G-orbit of some rigid algebra $\mathcal{J}$. In [1] he proved that every component of Assoc_n has to carry an open subset of non-singular points which is either the orbit of a single rigid algebra or an infinite union of orbits of algebras which differ only in the multiplication of the radical. Flanigan has discovered that the second alternative does in fact occur: there is a component of Assoc_3 which consists entirely of the orbits of an infinite family of three-dimensional nilpotent algebras.

In [2], G. Mazzola has proved that the subvariety of commutative algebras inside AssocN_n is irreducible for $n \leq 6$ and it admits at least two irreducible components in dimension 7. For the non-commutative case the description is known only for $n \leq 5$. A. MakhLouf showed that AssocN_2 is irreducible, while the number of irreducible components of AssocN_n when n is equal to 3, 4 and 5 is two, four and thirteen, respectively. Further, the author estimated a lower bound of the number of irreducible components in any dimension, see [3].

For the variety of nilpotent Lie algebras LieN_n the geometric classification is known for $n \leq 8$. In 1988, F. Grunewald and J. O'Halloran in [4] established that LieN_n is irreducible when $n \leq 5$. Two years later, S. Craig proved that this is the case for dimension 6 as well, see [5]. The first reducible variety of nilpotent Lie algebras appears in dimension 7, see [6]. It was shown by M. Goze and J. M. Ancochea-Bermudez. that LieN_7 decomposes into two algebraic components each corresponding to infinite family of algebras. Finally, in [7] the authors described all eight components of the variety of nilpotent Lie algebras of dimension 8.

As for the variety of Jordan algebras the references are rather recent. In [8, 2005], [9, 2014] it was shown that in $\mathcal{J}\mathrm{or}_n$ for $n \leq 4$ any irreducible component is determined by a rigid algebra. In [10, 2011], the authors establish that the subvariety of nilpotent algebras $\mathcal{J}\mathrm{orN}_n$ is irreducible for $n \leq 3$ and admits two irreducible components if $n = 4$.

In this paper, we will show that $\mathcal{J}\mathrm{orN}_5$ is the union of five irreducible components, there are four rigid algebras and the closure of the union of G-orbits of a certain infinite family gives rise to the fifth component.

The paper is organized as follows. Section 2 begins with some preliminaries on Jordan algebras, further we establish the list of non-isomorphic nilpotent Jordan algebras of dimension 5. Among the non-isomorphic nilpotent Jordan algebras there are four infinite families depending on one or two parameters from $\mathbf{k}$. In Section 3, we define the variety of Jordan algebras $\mathcal{J}\mathrm{orN}_n$, determine useful invariants of deformation and show how to construct deformation between algebras or how to ensure that such deformation does not exist. In Section 4 we give several examples for the deformation both of and into the infinite families, while in Section 5, we prove the main result of this paper describing deformations between the algebras in $\mathcal{J}\mathrm{orN}_5$ and characterizing its the irreducible components.

2 Jordan algebras: definition and basic concepts, list of nilpotent algebras up to dimension five

In this section we present the basic concepts, notations for Jordan algebras and the algebraic classification of nilpotent Jordan algebras of small dimensions.

Definition 1. A **Jordan $\mathbf{k}$-algebra** is a commutative $\mathbf{k}$-algebra $\mathcal{J}$ satisfying the Jordan identity, namely

$$((x \cdot x) \cdot y) \cdot x = (x \cdot x) \cdot (y \cdot x), \qquad x, y \in \mathcal{J}. \tag{1}$$

In a Jordan algebra $\mathcal{J}$ we define inductively a series of subsets by setting

$$\mathcal{J}^1 = \mathcal{J},$$
$$\mathcal{J}^n = \mathcal{J}^{n-1} \cdot \mathcal{J} + \mathcal{J}^{n-2} \cdot \mathcal{J}^2 + \cdots + \mathcal{J} \cdot \mathcal{J}^{n-1}.$$

The subset $\mathcal{J}^n$ is called the n**-th power of the algebra** $\mathcal{J}$.

Definition 2. A Jordan algebra $\mathcal{J}$ is said to be **nilpotent** if there exists an integer $n \in \mathbb{N}$ such that $\mathcal{J}^n = 0$. The least such number is called the **index of nilpotency** of the algebra $\mathcal{J}$, $\mathrm{nilindex}(\mathcal{J})$.

Further in this section we establish a list of all non-isomorphic nilpotent algebras of dimension five. For convenience we first describe all indecomposable nilpotent algebras of dimension up to four, using the notations from [11]. Also we drop $\cdot$ and denote a multiplication in $\mathcal{J}$ simply as xy. Products that we do not specify are understood to be 0 or determined by commutativity.

Example 3 (*Indecomposable nilpotent Jordan algebras of dimension one, and two three)***.** All nilpotent Jordan algebras of dimension less than four are associative. There is only one nilpotent indecomposable algebra in one and two: $\mathbf{k}n$, with $n^2 = 0$ and $\mathcal{B}_3$ generated by n_1, with $n_1^3 = 0$, respectively. There are two indecomposable nilpotent algebras in dimension three, $\mathcal{T}_3$ generated by n_1, such that $n_1^4 = 0$ and $\mathcal{T}_4 = \langle n_1, n_2 \mid n_1^2 = n_2^2 \rangle$.

In [11] all Jordan algebras of dimension four over an algebraically closed field of characteristic not 2 were described. In the Table 1 we list eight non-isomorphic indecomposable nilpotent Jordan algebras.

Table 1: Indecomposable four-dimensional nilpotent Jordan algebras.

$\mathcal{F}$	Multiplication Table
$\mathcal{F}_{61}$	$n_1^2 = n_2 \quad n_2^2 = n_4 \quad n_1n_2 = n_3 \quad n_1n_3 = n_4$
$\mathcal{F}_{62}$	$n_1^2 = n_2 \quad n_4^2 = n_2 \quad n_1n_2 = n_3$
$\mathcal{F}_{63}$	$n_1^2 = n_2 \quad n_4^2 = -n_2 - n_3 \quad n_1n_2 = n_3 \quad n_2n_4 = n_3$
$\mathcal{F}_{64}$	$n_1^2 = n_2 \quad n_4^2 = -n_2 \quad n_1n_2 = n_3 \quad n_2n_4 = n_3$
$\mathcal{F}_{65}$	$n_1^2 = n_2 \quad n_1n_2 = n_3 \quad n_2n_4 = n_3$
$\mathcal{F}_{66}$	$n_1^2 = n_2 \quad n_4^2 = n_3 \quad n_1n_2 = n_3$
$\mathcal{F}_{69}$	$n_1^2 = n_2 \quad n_1n_3 = n_4$
$\mathcal{F}_{70}$	$n_1^2 = n_2 \quad n_3n_4 = n_2$

2.1. Nilpotent Jordan algebras of dimension five

In [2], G. Mazzola has classified nilpotent commutative associative (and thus Jordan) five-dimensional algebras. Keeping the notations from the paper we list the 25 non-isomorphic such algebras in Table 2. For each algebra we calculate the dimension of its automorphism group $\mathrm{Aut}(\mathcal{J})$, of its annihilator $\mathrm{Ann}(\mathcal{J}) = \{a \in \mathcal{J} \mid a\mathcal{J} = 0\}$, of its second power $\mathcal{J}^2$ and the index of nilpotency $\mathrm{nilindex}(\mathcal{J})$.

Table 2: Five-dimensional nilpotent associative Jordan algebras.

ϵ	Multiplication Table	dim Aut	dim Ann	dim ϵ^2	nil-index
ϵ_1	$n_1^2 = n_2 \quad n_2^2 = n_1n_3 = n_4$ $n_1n_2 = n_3 \quad n_1n_4 = n_2n_3 = n_5$	5	1	4	6
ϵ_2	$n_1^2 = n_2 \quad n_2^2 = n_1n_3 = n_4$ $n_1n_2 = n_3$	6	1	3	5
ϵ_3	$n_1^2 = n_2 \quad n_4^2 = n_5$ $n_1n_2 = n_4n_5 = n_3$	6	1	3	4
ϵ_4	$n_1^2 = n_3 \quad n_1n_2 = n_4$ $n_1n_4 = n_2n_3 = n_5$	7	1	3	4
ϵ_5	$\mathcal{F}_{61} \oplus \mathbf{k}n_5$	7	2	3	5
ϵ_6	$\mathcal{T}_3 \oplus \mathcal{B}_3$	7	2	3	4
ϵ_7	$n_1n_3 = n_4 \quad n_2n_3 = n_5$ $n_1n_2 = n_4 - n_5$	7	2	2	3
ϵ_8	$n_1^2 = n_3 \quad n_2^2 = n_1n_3 = n_4$ $n_1n_2 = n_5$	8	2	3	4
ϵ_9	$n_1n_3 = n_5$ $n_1n_2 = n_4 \quad n_2n_3 = -n_5$	8	2	2	3
ϵ_{10}	$n_1^2 = n_3 \quad n_1n_3 = n_4 \quad n_1n_2 = n_5$	9	2	3	4
ϵ_{11}	$n_1^2 = n_2n_3 = n_4 \quad n_1n_3 = n_5$	9	2	2	3
ϵ_{12}	$n_1n_2 = n_4 \quad n_1n_3 = n_5$	11	2	2	3
ϵ_{13}	$n_3^2 = n_4 \quad n_1n_2 = n_3n_4 = n_5$	8	1	2	4
ϵ_{14}	$\mathcal{F}_{66} \oplus \mathbf{k}n_5$	9	2	2	4
ϵ_{15}	$\mathcal{T}_4 \oplus \mathcal{B}_3$	9	2	2	3
ϵ_{16}	$n_1^2 = n_3 \quad n_2^2 = n_5 \quad n_1n_2 = n_4$	10	3	3	3
ϵ_{17}	$n_1^2 = n_4 \quad n_3^2 = n_1n_2 = n_5$	10	2	2	3
ϵ_{18}	$\mathcal{T}_3 \oplus \mathbf{k}n_4 \oplus \mathbf{k}n_5$	11	3	2	4
ϵ_{19}	$\mathcal{B}_3 \oplus \mathcal{B}_3 \oplus \mathbf{k}n_5$	11	3	2	3
ϵ_{20}	$\mathcal{F}_{69} \oplus \mathbf{k}n_5$	12	3	2	3
ϵ_{21}	$n_2n_3 = n_1n_4 = n_5$	11	1	1	3
ϵ_{22}	$\mathcal{F}_{70} \oplus \mathbf{k}n_5$	12	2	1	3
ϵ_{23}	$\mathcal{T}_4 \oplus \mathbf{k}n_4 \oplus \mathbf{k}n_5$	14	3	1	3
ϵ_{24}	$\mathcal{B}_3 \oplus \mathbf{k}n_3 \oplus \mathbf{k}n_4 \oplus \mathbf{k}n_5$	17	4	1	3
ϵ_{25}	$\mathbf{k}n_1 \oplus \mathbf{k}n_2 \oplus \mathbf{k}n_3 \oplus \mathbf{k}n_4 \oplus \mathbf{k}n_5$	25	5	0	2

Recently, in [12] a method to construct nilpotent Jordan algebras using central extension of lower dimension algebras was presented. The authors showed that any nilpotent algebra can be obtained via this methods and provided a list of 35 single non-associative Jordan algebras together with 6 families of algebras depending on parameters in $\mathbf{k}$. The list of these algebras is given in Table 3 (using the notations [12]) together with the dimensions of $\mathrm{Aut}(\mathcal{J})$, $\mathrm{Ann}(\mathcal{J})$, $\mathcal{J}^2$ and of the associative center $\mathrm{Z}(\mathcal{J}) = \{a \in \mathcal{J} \mid (a, \mathcal{J}, \mathcal{J}) = (\mathcal{J}, a, \mathcal{J}) = (\mathcal{J}, \mathcal{J}, a) = 0\}$.

Table 3: Five-dimensional nilpotent non-associative Jordan algebras.

$\mathcal{J}$	Multiplication Table	dim Aut	dim Ann	dim $\mathcal{J}^2$	dim Z
$\mathcal{J}_1$	$\mathcal{F}_{65} \oplus \mathbf{k}n_5$	8	2	2	3
$\mathcal{J}_2$	$\mathcal{F}_{64} \oplus \mathbf{k}n_5$	9	2	2	3
$\mathcal{J}_3$	$\mathcal{F}_{63} \oplus \mathbf{k}n_5$	8	2	2	3
$\mathcal{J}_4$	$\mathcal{F}_{62} \oplus \mathbf{k}n_5$	7	2	2	3
$\mathcal{J}_5$	$n_1^2 = n_2 \quad n_4^2 = n_2n_3 = n_5$	6	1	2	3
$\mathcal{J}_6$	$n_1^2 = n_2 \quad n_1n_4 = n_2n_3 = n_5$	7	1	2	3
$\mathcal{J}_7$	$n_1n_2 = n_3 \quad n_3n_4 = n_5$	6	1	2	2
$\mathcal{J}_8$	$n_1n_2 = n_3 \quad n_3n_4 = n_1^2 = n_5$	5	1	2	2
$\mathcal{J}_9$	$n_1n_2 = n_3$ $n_3n_4 = n_1^2 = n_2^2 = n_5$	4	1	2	2
$\mathcal{J}_{10}$	$n_1n_2 = n_3$ $n_1n_3 = n_4^2 = n_2n_3 = n_5$	5	1	2	3
$\mathcal{J}_{11}$	$n_1n_2 = n_3 \quad n_1n_3 = n_4^2 = n_5$	7	1	2	3
$\mathcal{J}_{12}$	$n_1n_2 = n_3$ $n_1n_3 = n_4^2 = n_2^2 = n_5$	6	1	2	3
$\mathcal{J}_{13}$	$n_1n_2 = n_3$ $n_1n_4 = n_1n_3 = n_2n_3 = n_5$	6	1	2	3
$\mathcal{J}_{14}$	$n_1n_2 = n_3 \quad n_1n_3 = n_2n_4 = n_5$	7	1	2	3
$\mathcal{J}_{15}^{\alpha}$	$n_1^2 = n_2 \quad n_1n_2 = n_3 \quad n_4^2 = \alpha n_5$ $n_2^2 = n_1n_3 = n_2n_4 = n_5,\ \alpha \in \mathbf{k}$	6	1	3	3
$\mathcal{J}_{16}$	$n_1n_2 = n_3^2 = n_4 \quad n_1n_4 = n_5$	6	1	2	2
$\mathcal{J}_{17}$	$n_1n_2 = n_3^2 = n_4$ $n_1n_4 = n_2n_3 = n_5$	5	1	2	2
$\mathcal{J}_{18}$	$n_1n_2 = n_3^2 = n_4$ $n_1n_4 = n_2^2 = n_5$	4	1	2	2
$\mathcal{J}_{19}$	$n_1n_2 = n_3^2 = n_4 \quad n_3n_4 = n_5$	5	1	2	2
$\mathcal{J}_{20}$	$n_1n_2 = n_3^2 = n_4 \quad n_3n_4 = n_1^2 = n_5$	4	1	2	2
$\mathcal{J}_{21}$	$n_1n_2 = n_3^2 = n_4$ $n_3n_4 = n_1^2 = n_2^2 = n_5$	3	1	2	2
$\mathcal{J}_{22}$	$n_1^2 = n_2 \quad n_1n_2 = n_3^2 = n_4$ $n_2^2 = n_1n_4 = n_5$	4	1	3	3
$\mathcal{J}_{23}^{\beta}$	$n_1^2 = n_3 \quad n_1n_3 = n_2n_4 = n_5$ $n_1n_2 = n_4 \quad n_2n_3 = \beta n_5,\ \beta \in \mathbf{k}$	6	1	3	3
$\mathcal{J}_{24}^{\gamma}$	$n_1^2 = n_3 \quad n_1n_2 = n_4 \quad n_2n_3 = n_5$ $n_1n_4 = \gamma n_5,\ \gamma \in \mathbf{k}^* - \{1, -\frac{1}{2}\}$	7	1	3	3
$\mathcal{J}_{24}^{0}$	$n_1^2 = n_3 \quad n_1n_2 = n_4 \quad n_2n_3 = n_5$	7	2	3	3
$\mathcal{J}_{24}^{-\frac{1}{2}}$	$n_1^2 = n_3 \quad n_1n_2 = n_4 \quad n_2n_3 = n_5$ $n_1n_4 = -\frac{1}{2}n_5$	8	1	3	3
$\mathcal{J}_{25}$	$n_1^2 = n_3 \quad n_1n_2 = n_4$ $n_1n_3 = n_1n_4 = n_5 \quad n_2n_3 = -2n_5$	7	1	3	3
$\mathcal{J}_{26}^{\delta}$	$n_1^2 = n_3 \quad n_2^2 = n_4 \quad n_2n_4 = \delta n_5$ $n_1n_3 = n_1n_4 = n_5,\ \delta \in \mathbf{k}^*$	6	1	3	3

Table 3: Five-dimensional nilpotent non-associative Jordan algebras.

$\mathcal{J}$	Multiplication Table	dim Aut	dim Ann	dim $\mathcal{J}^2$	dim Z
$\mathcal{J}_{26}^{0}$	$n_1^2 = n_3 \quad n_2^2 = n_4$ $n_1n_3 = n_1n_4 = n_5$	6	2	3	3
$\mathcal{J}_{27}^{\varepsilon,\phi}$	$n_1^2 = n_3 \quad n_2n_3 = n_1n_4 = n_5$ $n_2^2 = n_4 \quad n_2n_4 = \varepsilon n_5$ $n_1n_3 = \phi n_5$ $\varepsilon,\phi\in\mathbf{k}$, $\varepsilon\phi\neq 1$	6	1	3	3
$\mathcal{J}_{27}^{\varepsilon,\varepsilon^{-1}}$	$n_1^2 = n_3 \quad n_2n_3 = n_1n_4 = n_5$ $n_2^2 = n_4 \quad n_2n_4 = \varepsilon n_5$ $n_1n_3 = \varepsilon^{-1}n_5 \quad \varepsilon\in\mathbf{k}^*-\{-1\}$	6	2	3	3
$\mathcal{J}_{27}^{-1,-1}$	$n_1^2 = n_3 \quad n_2n_3 = n_1n_4 = n_5$ $n_2^2 = n_4 \quad n_2n_4 = n_1n_3 = -n_5$	6	2	3	3
$\mathcal{J}_{28}$	$n_1^2 = n_2 \quad n_2n_3 = n_4 \quad n_3^2 = n_5$	7	2	3	3
$\mathcal{J}_{29}$	$n_1^2 = n_2 \quad n_2n_3 = n_4 \quad n_1n_3 = n_5$	7	2	3	3
$\mathcal{J}_{30}$	$n_1^2 = n_2 \quad n_2n_3 = n_4$ $n_1n_3 = n_3^2 = n_5$	6	2	3	3
$\mathcal{J}_{31}$	$n_1^2 = n_2 \quad n_2n_3 = n_4 \quad n_1n_2 = n_5$	7	2	3	3
$\mathcal{J}_{32}$	$n_1^2 = n_2 \quad n_2n_3 = n_4$ $n_1n_2 = n_1n_3 = n_5$	6	2	3	3
$\mathcal{J}_{33}$	$n_1^2 = n_2 \quad n_2n_3 = n_4$ $n_1n_2 = n_3^2 = n_5$	5	2	3	3
$\mathcal{J}_{34}$	$n_1n_2 = n_3 \quad n_1n_3 = n_4 \quad n_1^2 = n_5$	8	2	3	3
$\mathcal{J}_{35}$	$n_1n_2 = n_3 \quad n_1n_3 = n_4 \quad n_2^2 = n_5$	7	2	3	3
$\mathcal{J}_{36}$	$n_1n_2 = n_3 \quad n_1n_3 = n_4$ $n_2n_3 = n_5$	6	2	3	3
$\mathcal{J}_{37}$	$n_1^2 = n_2^2 = n_5 \quad n_1n_2 = n_3$ $n_1n_3 = n_4$	6	2	3	3
$\mathcal{J}_{38}$	$n_1n_2 = n_3 \quad n_1n_3 = n_4$ $n_1^2 = n_2n_3 = n_5$	5	2	3	3
$\mathcal{J}_{39}$	$n_1n_2 = n_3 \quad n_1^2 = n_5$ $n_2^2 = n_1n_3 = n_4$	7	2	3	3
$\mathcal{J}_{40}$	$n_1n_2 = n_3 \quad n_2^2 = n_1n_3 = n_4$ $n_1^2 = n_2n_3 = n_5$	4	2	3	3
$\mathcal{J}_{41}^{\lambda}$	$n_1^2 = n_5 \quad n_1n_2 = n_3 \quad n_2^2 = \lambda n_5$ $n_1n_3 = n_2n_3 = n_4,\ \lambda\in\mathbf{k}$	6	2	3	3

Theorem 4. *In Tables 2 and 3 all algebras are one-to-one non-isomorphic except for*

1. $\mathcal{J}_{15}^{\alpha_1} \simeq \mathcal{J}_{15}^{\alpha_2}$, $\mathcal{J}_{24}^{\alpha_1} \simeq \mathcal{J}_{24}^{\alpha_2}$ *if and only if* $\alpha_1 = \alpha_2$;
2. $\mathcal{J}_{23}^{\beta_1} \simeq \mathcal{J}_{23}^{\beta_2}$, $\mathcal{J}_{26}^{\beta_1} \simeq \mathcal{J}_{26}^{\beta_2}$ *if and only if* $\beta_1 = \pm\beta_2$;
3. $\mathcal{J}_{27}^{\alpha_1,\beta_1} \simeq \mathcal{J}_{27}^{\alpha_2,\beta_2}$ *if and only if* $(\alpha_1,\beta_1) = (\alpha_2,\beta_2)$ *or* $(\alpha_1,\beta_1) = (\beta_2,\alpha_2)$;
4. $\mathcal{J}_{41}^{\lambda_1} \simeq \mathcal{J}_{41}^{\lambda_2}$ *if and only if* $\lambda_1 = \lambda_2$ *or* $\lambda_1 = \lambda_2^{-1}$;
5. $\mathcal{J}_{24}^{1}$ *is in fact associative and corresponds to* ϵ_4 *from Table 2;*

6. $\mathcal{J}_{29} \simeq \mathcal{J}_{24}^{0}$;

7. $\mathcal{J}_{37} \simeq \mathcal{J}_{27}^{-1,-1}$.

8. *The family* $\mathcal{J}_{41}^{\lambda}$ *is isomorphic to* $\mathcal{J}_{27}^{\varepsilon,\varepsilon^{-1}}$ *for* $\lambda \neq 0,1$ *and* $\varepsilon = -\frac{(-1+\sqrt{\lambda})^2}{(1+\sqrt{\lambda})^2}$, *while* $\mathcal{J}_{41}^{1}$ *is isomorphic to* $\mathcal{J}_{26}^{0}$. *Remaining only* $\mathcal{J}_{41}^{0}$, *which will be denoted simply as* $\mathcal{J}_{41}$.

Proof. Firstly, in [12] the authors showed the isomorphism from items 1-4. The isomorphism 5 follows immediately from the multiplication tables of the corresponding algebras. To see 6-8 consider the change of basis

$\mathcal{J}_{29} \simeq \mathcal{J}_{24}^{0}$	$\mathcal{J}_{37} \simeq \mathcal{J}_{27}^{-1,-1}$	$\mathcal{J}_{41}^{1} \simeq \mathcal{J}_{26}^{0}$
$e_1 = n_1$	$e_1 = n_1 - n_2$	$e_1 = n_1 + n_2$
$e_2 = n_3$	$e_2 = -n_1 - n_2$	$e_2 = in_1 - in_2$
$e_3 = n_2$	$e_3 = -2(n_3 - n_5)$	$e_3 = 2(n_3 + n_5)$
$e_4 = n_5$	$e_4 = 2(n_3 + n_5)$	$e_4 = 2(n_3 - n_5)$
$e_5 = n_4$	$e_5 = 2n_4$	$e_5 = 4n_4$

and for any $\lambda \neq 0,1$ consider $\varepsilon = -\frac{(-1+\sqrt{\lambda})^2}{(1+\sqrt{\lambda})^2}$ then the transformation

$$
\begin{aligned}
e_1 &= \sqrt{\lambda} n_1 + n_2 \\
e_2 &= \left(\frac{\sqrt{\lambda} - \lambda}{1+\sqrt{\lambda}}\right) n_1 + \left(\frac{-1+\sqrt{\lambda}}{1+\sqrt{\lambda}}\right) n_2 \\
e_3 &= 2\sqrt{\lambda} n_3 + 2\lambda n_5 \\
e_4 &= -\frac{2(-1+\sqrt{\lambda})^2\sqrt{\lambda}}{(1+\sqrt{\lambda})^2} n_3 + \frac{2(-1+\sqrt{\lambda})^2\lambda}{(1+\sqrt{\lambda})^2} n_5 \\
e_5 &= -\frac{2(-1+\sqrt{\lambda})^2\sqrt{\lambda}}{1+\sqrt{\lambda}} n_4
\end{aligned}
$$

gives the isomorphism $\mathcal{J}_{41}^{\lambda} \simeq \mathcal{J}_{27}^{\varepsilon,\varepsilon^{-1}}$.

Further the dimension of the automorphism group, the annihilator, any power of algebra, the associative center, index of nilpotency together with associative and non-associative conditions, are invariants of class of isomorphic algebras. Therefore any two algebras whose last four columns in Table 2 or Table 3 are not equal are non-isomorphic, and one has to verify that the algebras are non-isomorphic in the following cases:

1. $\mathcal{J}_9 - \mathcal{J}_{18} - \mathcal{J}_{20}$: the four-dimensional non-associative nilpotent Jordan algebra $\mathcal{F}_{64}$ is a subalgebra of $\mathcal{J}_{20}$ but it is not a subalgebra nor of $\mathcal{J}_9$ neither of $\mathcal{J}_{18}$. Analogously, $\mathcal{F}_{70}$ is a subalgebra of $\mathcal{J}_9$ and is not of $\mathcal{J}_{18}$.

2. $\mathcal{J}_8 - \mathcal{J}_{17} - \mathcal{J}_{19}$: the dimension of the second cohomology group of $\mathcal{J}_{19}$ with coefficients in $\mathcal{J}_{19}$, $H^2(\mathcal{J}_{19}, \mathcal{J}_{19})$, is 6 while $\dim H^2(\mathcal{J}_8, \mathcal{J}_8) = \dim H^2(\mathcal{J}_{17}, \mathcal{J}_{17}) = 5$. For the definition of the cohomology group for Jordan algebras we refer to [9]. On the other hand, the four-dimensional algebra $\mathcal{F}_{63}$ is a subalgebra of $\mathcal{J}_8$ but is not of $\mathcal{J}_{17}$.

3. $\mathcal{J}_{33} \not\simeq \mathcal{J}_{38}$: the three-dimensional nilpotent Jordan algebra $\mathcal{T}_4$ is a subalgebra of $\mathcal{J}_{38}$ but is not of $\mathcal{J}_{33}$.

4. $\mathcal{J}_5 - \mathcal{J}_{12} - \mathcal{J}_{13}$: $\dim H^2(\mathcal{J}_{12}, \mathcal{J}_{12}) = \dim H^2(\mathcal{J}_{13}, \mathcal{J}_{13}) = 13$ while $\dim H^2(\mathcal{J}_5, \mathcal{J}_5) = 12$ But the dimension of Jacobi space of $\mathcal{J}_{12}$, i.e.

$$\mathrm{Jac}(\mathcal{J}) = \{a \in \mathcal{J} \mid a(xy) = (ax)y + x(ay),\ \forall x, y \in \mathcal{J}\},$$

is 4 whereas $\dim \mathrm{Jac}(\mathcal{J}_{13}) = 3$.

5. $\mathcal{J}_7 \not\simeq \mathcal{J}_{16}$: the four-dimensional algebra $\mathcal{F}_{71} = \mathcal{T}_4 \oplus \mathbf{k}n_4$ is a subalgebra of $\mathcal{J}_7$ but is not of $\mathcal{J}_{16}$.

6. $\mathcal{J}_6 - \mathcal{J}_{11} - \mathcal{J}_{14}$: $\mathcal{F}_{63} \subseteq \mathcal{J}_{14}$ while $\mathcal{F}_{63} \not\subseteq \mathcal{J}_6$, adding $\dim H^2(\mathcal{J}_{11}, \mathcal{J}_{11}) = 15$ while $\dim H^2(\mathcal{J}_6, \mathcal{J}_6) = \dim H^2(\mathcal{J}_{14}, \mathcal{J}_{14}) = 14$ we obtain the result.

7. $\mathcal{J}_{24}^{\gamma} \not\simeq \mathcal{J}_{25}$: for any $\gamma \neq 0$, algebra $\mathcal{F}_{69}$ is a subalgebra of $\mathcal{J}_{24}^{\gamma}$ but is not a subalgebra of $\mathcal{J}_{25}$. Since $\dim \mathrm{Ann}(\mathcal{J}_{24}^0) = 2$, $\mathcal{J}_{25} \not\simeq \mathcal{J}_{24}^0$.

8. $\mathcal{J}_{28} - \mathcal{J}_{24}^0 - \mathcal{J}_{31} - \mathcal{J}_{35} - \mathcal{J}_{39}$: one compares the dimensions of both second cohomology group and Jacobi space, $\dim H^2(\mathcal{J}_{35}, \mathcal{J}_{35}) = 10$, $\dim H^2(\mathcal{J}_{31}, \mathcal{J}_{31}) = 8$, $\dim H^2(\mathcal{J}_{28}, \mathcal{J}_{28}) = \dim H^2(\mathcal{J}_{39}, \mathcal{J}_{39}) = 11$, and $\dim H^2(\mathcal{J}_{24}^0, \mathcal{J}_{24}^0) = 12$, while $\dim \mathrm{Jac}(\mathcal{J}_{28}) = 3$ and $\dim \mathrm{Jac}(\mathcal{J}_{39}) = 4$.

9. $\mathcal{J}_{15}^{\alpha} - \mathcal{J}_{23}^{\beta} - \mathcal{J}_{26}^{\delta} - \mathcal{J}_{27}^{\varepsilon,\phi}$: the $\mathrm{nilindex}(\mathcal{J}_{15}^{\alpha}) = 5$ for any $\alpha \in \mathbf{k}$ while it is 4 for the other algebras. Analogously, $\dim H^2(\mathcal{J}_{23}^{\beta}, \mathcal{J}_{23}^{\beta}) = 8$ if $\beta \neq 0$ and $\dim H^2(\mathcal{J}_{23}^0, \mathcal{J}_{23}^0) = 9$, while $\dim H^2(\mathcal{J}_{26}^{\delta}, \mathcal{J}_{26}^{\delta}) = \dim H^2(\mathcal{J}_{27}^{\varepsilon,\phi}, \mathcal{J}_{27}^{\varepsilon,\phi}) = 7$ if $\delta \neq 0$ and $\varepsilon\phi \neq 1$.

   Next, suppose that $\mathcal{J}_{27}^{\varepsilon,\phi} \simeq \mathcal{J}_{26}^{\delta}$ for some $\delta, \varepsilon, \phi \in \mathbf{k}$ and suppose that this isomorphism is given by a matrix $A = (a_{ij})_{i,j=1}^5$. Then the coefficients a_{ij} have to satisfy the following conditions: $a_{ij} = a_{i1}a_{i2} = a_{k5} = 0$ for $i = 1, 2$, $j = 3, 4, 5$, and $a_{3k} = a_{1(k-2)}^2$, $a_{4k} = a_{2(k-2)}^2$, $a_{5k} = +2a_{2(k-2)}(a_{3(k-2)} + a_{4(k-2)}\varepsilon) + 2a_{1(k-2)}(a_{4(k-2)} + a_{3(k-2)}\phi) = 0$, for $k = 3, 4$. Moreover,

$$\begin{aligned}
a_{21}a_{32} + a_{11}a_{42} + a_{21}a_{42}\varepsilon + a_{41}(a_{12} + a_{22}\varepsilon) + a_{11}a_{32}\phi + a_{31}(a_{22} + a_{12}\phi) &= 0\\
a_{11}a_{21}(a_{11} + a_{21}) - a_{55} + a_{21}^3\varepsilon + a_{11}^3\phi &= 0\\
-a_{55} + a_{21}(a_{12}^2 + a_{22}^2\varepsilon) + a_{11}(a_{22}^2 + a_{12}^2\phi) &= 0\\
a_{21}^2(a_{12} + a_{22}\varepsilon) + a_{11}^2(a_{22} + a_{12}\phi) &= 0\\
a_{12}a_{22}(a_{12} + a_{22}) - a_{55}\delta + a_{22}^3\varepsilon + a_{12}^3\phi &= 0.
\end{aligned}$$

   But the system allows only trivial solutions $a_{i5} = 0$ for $1 \leq i \leq 5$.

10. $\mathcal{J}_{26}^0 - \mathcal{J}_{27}^{\varepsilon,\varepsilon^{-1}} - \mathcal{J}_{30} - \mathcal{J}_{32} - \mathcal{J}_{36} - \mathcal{J}_{27}^{-1,-1} - \mathcal{J}_{41}$: the nilpotency type of $\mathcal{J}_{32}$ and $\mathcal{J}_{36}$ is $(2, 1, 2)$ while for all other algebras it is $(2, 2, 1)$ (for definition of nilpotency type of Jordan algebra we refer to [11]). Suppose that $\mathcal{J}_{36} \simeq \mathcal{J}_{32}$ and the isomorphism is given by the matrix $A = (a_{ij})_{i,j=1}^5$. Then a_{ij}'s have to satisfy the following conditions: $a_{ij} = a_{3k} = 2a_{i3}a_{33} = 0$ for $i = 1, 2$, $j = 2, 4, 5$, $k = 4, 5$, $a_{n2} = 2a_{11}a_{(n-1)1}$, for $n = 3, 4$, and $a_{52} = 2a_{31}a_{21}$, $2a_{13}a_{23} = 0$, $2a_{11}a_{21}a_{23} = a_{54}$, $2a_{11}a_{21}a_{13} = a_{44}$, $a_{31}a_{23} + a_{21}a_{33} = a_{55} = 2a_{11}a_{21}^2$, $a_{31}a_{13} + a_{11}a_{33} = a_{45} = 2a_{11}^2a_{21}$, $a_{21}a_{13} + a_{11}a_{23} = 0$. This implies that $a_{i4} = 0$ for $1 \leq i \leq 5$.

    Comparing the dimensions of the cohomologies we rule out some cases:

$$\begin{gathered}
\dim H^2(\mathcal{J}_{27}^{-1,-1}, \mathcal{J}_{27}^{-1,-1}) = \dim H^2(\mathcal{J}_{41}, \mathcal{J}_{41}) = 8,\\
\dim H^2(\mathcal{J}_{27}^{\varepsilon,\varepsilon^{-1}}, \mathcal{J}_{27}^{\varepsilon,\varepsilon^{-1}}) = \dim H^2(\mathcal{J}_{30}, \mathcal{J}_{30}) = 7\\
\dim H^2(\mathcal{J}_{26}^0, \mathcal{J}_{26}^0) = 10.
\end{gathered}$$

On the other hand, $\dim \mathrm{Jac}(\mathcal{J}_{27}^{-1,-1}) = 4$ whereas $\dim \mathrm{Jac}(\mathcal{J}_{41}) = 3$. It remains to check that $\mathcal{J}_{27}^{\varepsilon,\varepsilon^{-1}} \not\simeq \mathcal{J}_{30}$. Suppose that $A = (a_{ij})_{i,j=1}^{5}$ which gives an isomorphism from $\mathcal{J}_{30}$ to $\mathcal{J}_{27}^{\varepsilon,\varepsilon^{-1}}$, then $a_{ij} = a_{k5} = 0$ for $i = 1, 3$, $j = 3, 4, 5$ and $k = 2, 5$. Moreover, $a_{11}^2 = a_{23}$, $2a_{21}a_{31} = a_{43}$, $2a_{11}a_{31} + a_{31}^2 = a_{53}$, $a_{11}a_{12} = 0$, $a_{31}a_{22} + a_{21}a_{32} = 0$, $a_{11}a_{32} + a_{31}(a_{12} + a_{32}) = 0$, $a_{32}a_{24}\varepsilon^{-1} = a_{31}a_{23}\varepsilon = a_{45} = a_{31}a_{24} = a_{32}a_{23}$, $a_{12}^2 = a_{24}$, $2a_{22}a_{32} = a_{44}$ and $2a_{12}a_{32} + a_{32}^2 = a_{54}$. All the solutions lead to $a_{i5} = 0$ for $1 \leq i \leq 5$, a contradiction.

□

3 Deformation, variety of Jordan algebras, constructing deformations

Let V be a n-dimensional $\mathbf{k}$-vector space with a fixed basis $\{e_1, e_2, \cdots, e_n\}$. To endow V with a Jordan $\mathbf{k}$-algebra structure, $(\mathcal{J}, \cdot)$, it suffices to specify n^3 structure constants $c_{ij}^k \in \mathbf{k}$, namely, $e_i \cdot e_j = \sum_{k=1}^{n} c_{ij}^k e_k$, $i, j \in \{1, 2, \ldots, n\}$. If a point $(c_{ij}^k) \in \mathbf{k}^{n^3}$ defines Jordan algebra its coordinates satisfy certain polynomial equations which follow from commutativity and Jordan identity. These equations cut out an algebraic variety $\mathcal{J}\mathrm{or}_n$ in $\mathbf{k}^{n^3}$. The points $(c_{ij}^k) \in \mathcal{J}\mathrm{or}_n$ are in one-to-one correspondence with n-dimensional $\mathbf{k}$-algebras along with a particular choice of basis. A change of basis in $\mathcal{J}$ is given by the general linear group $\mathrm{G} = \mathrm{GL}(\mathrm{V})$ action on $\mathcal{J}\mathrm{or}_n$, namely:

$$g(\mathcal{J}, \cdot) \;\mapsto\; (\mathcal{J}, \cdot_g), \qquad x \cdot_g y = g(g^{-1}x \cdot g^{-1}y), \tag{2}$$

for any $\mathcal{J} \in \mathcal{J}\mathrm{or}_n$, $g \in \mathrm{G}$ and $x, y \in \mathrm{V}$. The G-**orbit** of a Jordan algebra $\mathcal{J}$, that is, the set of all images of $\mathcal{J}$ under the action of G, is denoted by $\mathcal{J}^{\mathrm{G}}$. The set of different G-orbits of this action is in one-to-one correspondence with the set of isomorphism classes of n-dimensional Jordan algebras. The Zariski topology of the affine space provides the following relation of orbits, namely, we say that $\mathcal{J}_1$ is a **deformation** of $\mathcal{J}_2$ or that $\mathcal{J}_1$ **dominates** $\mathcal{J}_2$, if the orbit $\mathcal{J}_2^{\mathrm{G}}$ is contained in the Zariski closure of the orbit $\mathcal{J}_1^{\mathrm{G}}$. We denote it by $\mathcal{J}_1 \to \mathcal{J}_2$. A Jordan algebra $\mathcal{J}$ is called **rigid** if its G-orbit is a Zariski-open set in $\mathcal{J}\mathrm{or}_n$. Rigid algebra $\mathcal{J}$ does not admit non-trivial deformations, indeed if $\mathcal{J}_1 \to \mathcal{J}$ then $\mathcal{J}_1^{\mathrm{G}} \cap \mathcal{J}^{\mathrm{G}} \neq \varnothing$ and therefore $\mathcal{J}_1 \simeq \mathcal{J}$.

As we have already mentioned in the introduction the rigid algebras are of particular interest. Recall that any affine variety could be decomposed into its irreducible components. When $\mathcal{J} \in \mathcal{J}\mathrm{or}_n$ is rigid, there exists an irreducible component $\mathcal{C}$ such that $\mathcal{J}^{\mathrm{G}} \cap \mathcal{C}$ is a non-empty open subspace in $\mathcal{C}$, and therefore the closure of $\mathcal{J}^{\mathrm{G}}$ coincides with $\mathcal{C}$. The converse is not true: in the sections 5 we will show that there exists an irreducible component which does not contain a single open G-orbit but rather an open subset which consists of infinite union of orbits of algebras.

Any $(c_{ij}^k) \in \mathcal{J}\mathrm{or}_n$ which corresponds to a nilpotent Jordan algebra has to satisfy polynomial equation, thus the set of nilpotent Jordan algebras in dimension n possess a structure of algebraic variety embedded in $\mathcal{J}\mathrm{or}\mathrm{N}_n \subseteq \mathcal{J}\mathrm{or}_n$. Further in this section we collect the properties of deformation which will be used to describe $\mathcal{J}\mathrm{or}\mathrm{N}_5$. The first fact is straightforward but very useful.

Lemma 5. *[9] Let $\mathcal{J}_i$ and $\mathcal{J}_i'$ be in $\mathcal{J}\mathrm{or}_{n_i}$, for $i = 1, 2$. If $\mathcal{J}_i \to \mathcal{J}_i'$, then $\mathcal{J}_1 \oplus \mathcal{J}_2 \to \mathcal{J}_1' \oplus \mathcal{J}_2'$.*

Example 6. From the description of $\mathcal{J}\mathrm{or}\mathrm{N}_4$ in [10] it follows that $\mathcal{F}_{62} \to \mathcal{F}_{63} \to \mathcal{F}_{64}$ and $\mathcal{F}_{62} \to \mathcal{F}_{65}$, consequently $\mathcal{J}_4 \to \mathcal{J}_3 \to \mathcal{J}_2$ and $\mathcal{J}_4 \to \mathcal{J}_1$.

The other way to deform one algebra into another is to build a curve as specified in Lemma 7.

Lemma 7. *[2]Suppose that there exists a curve γ in $\mathcal{J}\mathrm{or}_n$ which generically lies in a subvariety $\mathcal{U}$ and which cuts $\mathcal{J}^{\mathrm{G}}$ in special point then $\mathcal{J} \in \overline{\mathcal{U}}$.*

Example 8. Let $\gamma(t) = \left\{e_1^t, e_2^t, e_3^t, e_4^t, e_5^t\right\}$ be a change of basis of $\mathcal{J}_{21}$ given by: $e_1^t = \sqrt{2}t^2 n_1 - \sqrt{2}t^2 n_2 + 2t^2 n_3 - t^2 n_4$, $e_2^t = 2\sqrt{2}t^4 n_2 + 4t^4 n_4$, $e_3^t = \sqrt{2}t^3(n_1 + n_2)$, $e_4^t = 4t^6(n_4 + n_5)$ and $e_5^t = 8t^8 n_5$. Thus γ defines a curve in $\mathcal{J}\mathrm{orN}_5$ which for any $t \neq 0$ belongs to $\mathcal{J}_{21}^{\mathrm{G}}$ (observe that $\det \gamma(t) = -2^8 t^{23}$). However, $e_2^t e_3^t = t e_4^t$ and for $t \to 0$ we get the multiplication table of $\mathcal{J}_{18}$. Thus, from Lemma 7, it follows that $\mathcal{J}_{18} \in \overline{\mathcal{J}_{21}^{\mathrm{G}}}$, i.e., $\mathcal{J}_{21} \to \mathcal{J}_{18}$.

The non-existence of deformation for a given pair of algebras $\mathcal{J}_1, \mathcal{J}_2 \in \mathcal{J}\mathrm{or}_n$ will follow from the violation of one of the conditions below.

Theorem 9. *If $\mathcal{J}_1 \to \mathcal{J}_2$ then:*

1. $\dim \mathrm{Aut}(\mathcal{J}_1) < \dim \mathrm{Aut}(\mathcal{J}_2)$*;*
2. $\dim \mathrm{Ann}(\mathcal{J}_1) \leq \dim \mathrm{Ann}(\mathcal{J}_2)$*;*
3. $\dim(\mathcal{J}_1^m) \geq \dim(\mathcal{J}_2^m)$*, for any positive integer m;*
4. $\dim \mathrm{Z}(\mathcal{J}_1) \leq \dim \mathrm{Z}(\mathcal{J}_2)$.

Proof. For the proof of (1) to (3) we refer to [9]. To see (4), it suffices to prove that the sets $\{\mathcal{J} \in \mathcal{J}\mathrm{or}_n \mid \dim \mathrm{Z}(\mathcal{J}) \geq s\}$ are Zariski-closed in $\mathcal{J}\mathrm{or}_n$ for any $s \in \mathbb{N}$. Let $e_1, e_2, \ldots, e_n$ be a basis for $\mathcal{J} \in \mathcal{J}\mathrm{or}_n$ and $a = \sum_{i=1}^n \alpha_i e_i \in \mathrm{Z}(\mathcal{J})$. Substituting a in each of the $3n^2$ equations $(a, e_i, e_j) = 0$, $(e_i, a, e_j) = 0$ and $(e_i, e_j, a) = 0$ for $i, j = 1, \ldots, n$, we obtain a system with $3n^3$ equations having α_i as unknowns. Then $\dim \mathrm{Z}(\mathcal{J}) = n - \mathrm{rank}(P_{3n^3,n})$, where $P_{3n^3,n}$ is the matrix of the system, thus the condition $\dim \mathrm{Z}(\mathcal{J}) \geq s$ is equivalent to the fact that all $(n - s + 1)$-minors of $P_{3n^3,n}$ vanish, which is defined by a finite number of polynomial identities in the structure constants. □

As direct consequences of (3) and (4), respectively, we have:

Corollary 10.

1. *If $\mathcal{J}_1$ and $\mathcal{J}_2$ are nilpotent Jordan algebras such as $\mathcal{J}_1 \to \mathcal{J}_2$ then* $\mathrm{nilindex}(\mathcal{J}_1) \geq \mathrm{nilindex}(\mathcal{J}_2)$.
2. *Any deformation of a non-associative algebra is a non-associative algebra.*

4 Deformations of infinite families of algebras

Whenever the class of algebras consists of only finitely many non-isomorphic algebras, the irreducible components are completely determined by rigid algebras, see $\mathcal{J}\mathrm{or}_n$, $n \leq 4$. When we have infinite number of non-isomorphic algebras it is not necessarily that any irreducible component contains an open orbit. In [1], Flanigan showed that there exists an infinite family of nilpotent algebras of dimension three such that the closure of a union of their orbits is an irreducible component in AssocN_3, while it carries no open orbit. In the next section, we will show that such irreducible component exists in $\mathcal{J}\mathrm{orN}_5$. In general, the behaviour of infinite families under deformation is of particular interest and in what

follows we will look into examples of deformations in $\mathcal{J}\mathrm{or}\mathrm{N}_5$ which involve families. First, to establish the notation denote by

$$\mathcal{N}_i^{\#} = \bigcup_{\alpha\in T} (\mathcal{J}_i^{\alpha})^{\mathrm{G}},$$

the union of the orbits of algebras in the family $\mathcal{J}_i^{\alpha}$, $\alpha \in T$ where T is some subset of $\mathbf{k}$.

4.1. Infinite family is dominated by a single algebra

The infinite family $\mathcal{J}_{15}^{\alpha},\ \alpha \in \mathbf{k}$, is dominated by algebra $\mathcal{J}_{22}$. To see that, for any $\alpha \in \mathbf{k}$ consider a change of basis of $\mathcal{J}_{22}$ parametrized by $\gamma_\alpha(t)$: $e_1^t = t^2n_1 - \frac{\alpha}{3}n_2 + tn_3$, $e_2^t = t^4n_2 + (1-\frac{2\alpha}{3})t^2n_4 + \frac{\alpha^2}{9}n_5$, $e_3^t = t^6n_4 - (-1+\alpha)t^4n_5$, $e_4^t = t^4n_2 - t^5n_3 + \frac{\alpha t^2}{3}n_4$ and $e_5^t = t^8n_5$. Observe that $\det(\gamma_\alpha) = t^{25}$ thus, for any $t \neq 0$, the curve $\gamma_\alpha(t)$ belongs to $\mathcal{J}_{22}^{\mathrm{G}}$. Since $\left(e_4^t\right)^2 = t^4e_3^t + \alpha e_5^t$ it follows that $\gamma_\alpha(0)$ gives the multiplication table of $\mathcal{J}_{15}^{\alpha}$. Therefore $\mathcal{N}_{15}^{\#} \subset \overline{\mathcal{J}_{22}^{\mathrm{G}}}$.

4.2. Infinite family is a deformation of a single algebra

The deformation from the family $\mathcal{J}_{23}^{\beta},\ \beta \in \mathbf{k}$ to $\mathcal{J}_{41}$ is quite remarkable, since no single member of $\mathcal{J}_{23}^{\beta}$ can dominate $\mathcal{J}_{41}$. Indeed $\dim \mathrm{Aut}(\mathcal{J}_{41}) = \dim \mathrm{Aut}(\mathcal{J}_{23}^{\beta})$ for all $\beta \in \mathbf{k}$. Consider the "variable" change of basis of $\mathcal{J}_{23}^{\beta(t)}$, where $\beta(t) = \frac{1}{t}$, namely: $e_1^t = tn_1 - \frac{1}{2}n_2$, $e_2^t = -\frac{1}{2}n_2$, $e_3^t = -\frac{t}{2}n_3$, $e_4^t = \frac{t}{4}n_5$ and $e_5^t = t^2n_3 - tn_4$. So the curve $\gamma(t)$ lies transversely to the orbits of $\mathcal{J}_{23}^{\beta}$, meaning that, for any $t \neq 0$, $\gamma(t) \subset \mathcal{N}_{23}^{\#}$ and cuts $\mathcal{J}_{41}^{\mathrm{G}}$ in $t = 0$. Thus $\mathcal{J}_{41} \in \overline{\mathcal{N}_{23}^{\#}}$. We will abuse the notation and will denote this fact by $\mathcal{J}_{23}^{\beta} \to \mathcal{J}_{41}$.

Observe that in the same way we obtain that any algebra $\mathcal{J}_k^{\alpha}$ of a family can be viewed as a deformation of $\mathcal{N}_k^{\#}$. We use the basis of $\mathcal{J}_k^{\alpha}$ from Table 3, then for any fixed α_0 $\gamma(\alpha)$ defines the multiplication in $\mathcal{J}_k^{\alpha_0}$ while for all other values of α algebra belongs to $\mathcal{N}_k^{\#}$.

In a rather different way, we obtain the dominance of $\mathcal{J}_{39}$ by the family $\mathcal{J}_{15}^{\alpha},\ \alpha \in \mathbf{k}$. In this case $\mathcal{J}_{15}^{1}$ (a single member of $\mathcal{J}_{15}^{\alpha},\ \alpha \in \mathbf{k}$) dominates $\mathcal{J}_{39}$. Consider the curve: $e_1^t = tn_1$, $e_2^t = n_2 + (-1+t)n_4$, $e_3^t = tn_3$, $e_4^t = t^2n_5$ and $e_5^t = t^2n_2$. For any $t \neq 0, 1$, the curve lies in $\left(\mathcal{J}_{15}^{1}\right)^{\mathrm{G}}$ and cuts $\mathcal{J}_{39}^{\mathrm{G}}$ in $t = 0$ and ϵ_5^{G} in $t = 1$, respectively. Then we have

$$\mathcal{J}_{39} \in \overline{\left(\mathcal{J}_{15}^{1}\right)^{\mathrm{G}}} \subseteq \bigcup_{\alpha\in\mathbf{k}} \overline{\left(\mathcal{J}_{15}^{\alpha}\right)^{\mathrm{G}}} \subseteq \overline{\mathcal{N}_{15}^{\#}}$$

and, therefore, $\mathcal{J}_{15}^{\alpha} \to \mathcal{J}_{39}$.

4.3. Deformations between families

The family $\mathcal{J}_{27}^{\varepsilon,\phi}$, $\varepsilon, \phi \in \mathbf{k}$ and $\varepsilon\phi \neq 1$, dominates the family $\mathcal{J}_{26}^{\delta}$, $\delta \in \mathbf{k}^*$. Again no single member of $\mathcal{J}_{26}^{\delta}$ deforms into any single member of $\mathcal{J}_{27}^{\varepsilon,\phi}$ since $\dim \mathrm{Aut}(\mathcal{J}_{26}^{\delta}) = \dim \mathrm{Aut}(\mathcal{J}_{27}^{\varepsilon,\phi})$, for all $\delta, \varepsilon, \phi \in \mathbf{k}$. For any fixed $\delta \in \mathbf{k}^*$, we construct a curve $\gamma_\delta(t)$ which lie transversely to the orbits of $\mathcal{J}_{27}^{\varepsilon,\phi}$. Consider the "variable" change of basis of $\mathcal{J}_{27}^{\varepsilon(t),\phi(t)}$, where $(\varepsilon(t), \phi(t)) = (t\delta, \frac{1}{t^2})$, defined as: $e_1^t = t\delta n_1$, $e_2^t = \delta n_2$, $e_3^t = t^2\delta^2 n_3$, $e_4^t = \delta^2 n_4$ and $e_5^t = t\delta^3 n_5$. For any $t \neq 0$, the curve lies in $\mathcal{N}_{27}^{\#}$ and, since $e_2^t e_3^t = te_5^t$, cuts $\left(\mathcal{J}_{26}^{\delta}\right)^{\mathrm{G}}$ in $t = 0$. Thus $\mathcal{J}_{26}^{\delta} \in \overline{\mathcal{N}_{27}^{\#}}$, for any $\delta \in \mathbf{k}^*$.

Differently, we obtain the dominance of the family $\mathcal{J}_{24}^{\gamma}$, $\gamma \in \mathbf{k}^* - \{1, -\frac{1}{2}\}$ by the family $\mathcal{J}_{23}^{\beta},\ \beta \in \mathbf{k}$. In this case, fixing $\gamma_0 \in \mathbf{k}^* - \{1, -\frac{1}{2}\}$, the algebra $\mathcal{J}_{24}^{\gamma_0}$ deforms into $\mathcal{J}_{23}^{\beta_0}$, where $\beta_0 = \frac{(1-2\gamma_0)i}{\sqrt{\gamma_0}}$. Define the curve: $e_1^t = -\frac{i}{\sqrt{\gamma_0}}n_1 + n_2$, $e_2^t = tn_2$, $e_3^t = -\frac{1}{\gamma_0}n_3 - \frac{2i}{\sqrt{\gamma_0}}n_4$,

$e_4^t = -\frac{it}{\sqrt{\gamma_0}} n_4$ and $e_5^t = -\frac{it}{\sqrt{\gamma_0^3}} n_5$. For any $t \neq 0$, the curve lies in the G-orbit of $\mathcal{J}_{23}^{\beta_0}$ and, since $e_2^t e_4^t = t\gamma_0 e_5^t$, cuts $\left(\mathcal{J}_{24}^{\gamma_0}\right)^{\mathrm{G}}$ in $t = 0$. Therefore

$$\mathcal{J}_{24}^{\gamma_0} \in \overline{\left(\mathcal{J}_{23}^{\beta_0}\right)^{\mathrm{G}}} \subseteq \overline{\mathcal{N}_{23}^{\#}}$$

consequently $\mathcal{J}_{23}^{\beta} \to \mathcal{J}_{24}^{\gamma}$, for all $\gamma \in \mathbf{k}^* - \{1, -\frac{1}{2}\}$.

5 The algebraic variety $\mathcal{J}\mathrm{or}\mathrm{N}_5$

In this section, we will describe the algebraic variety of five-dimensional Jordan algebras $\mathcal{J}\mathrm{or}\mathrm{N}_5$. First we show that any structure can be deformed either into one of the given four algebras or into the algebras in family $\mathcal{J}_{27}^{\varepsilon,\phi}$. The main tool of the proof is Lemma 7, to define a curve $\gamma(t)$ we use notations and basis of Table 3.

Lemma 11. *Every structure $\mathcal{J} \in \mathcal{J}\mathrm{or}\mathrm{N}_5$ belongs to $\overline{\epsilon_1^{\mathrm{G}}}$, $\overline{\mathcal{J}_{21}^{\mathrm{G}}}$, $\overline{\mathcal{J}_{22}^{\mathrm{G}}}$, $\overline{\mathcal{J}_{40}^{\mathrm{G}}}$ or $\overline{\mathcal{N}_{27}^{\#}}$.*

Proof. All associative structures, namely ϵ_i for $i = 1, \ldots, 25$, belong to $\overline{\epsilon_1^{\mathrm{G}}}$. In fact, G. Mazzola in [2] showed that the subvariety of associative algebras is an irreducible subvariety in $\mathcal{J}\mathrm{or}\mathrm{N}_5$ and that the orbit ϵ_1^{G} is open, thus its closure contains all associative structure. From Corollary 10 it follows that there is no non-associative structure which deforms into ϵ_1.

All algebras $\mathcal{J}_i$ for $i = 1, \ldots, 20$ except for the family $\mathcal{J}_{15}^{\alpha}$, are dominated by $\mathcal{J}_{21}$. Example 8 gives $\mathcal{J}_{21} \to \mathcal{J}_{18}$. For the deformation $\mathcal{J}_{18} \to \mathcal{J}_{17}$, consider a family of algebras with the following basis

$$\begin{array}{l} e_1^t = -t^2 n_1 + \frac{1}{2} n_2 - t n_3 + \frac{1}{8t^2} n_4, \\ e_2^t = -t^8 n_2 - \frac{3t^6}{2} n_4, \\ e_3^t = t^4 n_2 - t^5 n_3 + \frac{t^2}{2} n_4, \\ e_4^t = t^{10} n_4 + t^8 n_5, \\ e_5^t = -t^{12} n_5. \end{array} \tag{3}$$

Here $\{n_1, \ldots, n_5\}$ is the basis of $\mathcal{J}_{18}$ given in Table 3, $t \in \mathbf{k}$ and therefore this family can be viewed as a curve $\gamma(t) \in \mathcal{J}\mathrm{or}\mathrm{N}_5$. For any fixed $t \neq 0$ the change of basis (3) determines algebras isomorphic to $\mathcal{J}_{18}$, while for $t = 0$ one obtains $\mathcal{J}_{17}$ (indeed, $\left(e_2^t\right)^2 = -t^4 e_5^t$). Analogously, the change of basis $e_1^t = n_1$, $e_2^t = t^2 n_2$, $e_3^t = t n_3$, $e_4^t = t^2 n_4$ and $e_5^t = t^2 n_5$ of $\mathcal{J}_{17}$ does the trick for $\mathcal{J}_{17} \to \mathcal{J}_{16}$. Applying Lemma 7 to given transformations we obtain the following deformations:

$\mathcal{J}_{21} \to \mathcal{J}_{20}$	$\mathcal{J}_{20} \to \mathcal{J}_{19}$	$\mathcal{J}_{19} \to \mathcal{J}_{12}$	$\mathcal{J}_{12} \to \mathcal{J}_{11}$	$\mathcal{J}_{20} \to \mathcal{J}_{8}$
$e_1^t = \frac{1}{t^3} n_1$	$e_1^t = t^2 n_1$	$e_1^t = n_1 - \frac{1}{2} n_2 + n_3$	$e_1^t = n_1$	$e_1^t = t n_1$
$e_2^t = \frac{1}{t} n_2$	$e_2^t = n_2$	$e_2^t = t n_3 + n_4$	$e_2^t = t^2 n_2$	$e_2^t = n_2$
$e_3^t = \frac{1}{t^2} n_3$	$e_3^t = t n_3$	$e_3^t = t n_4 + n_5$	$e_3^t = t^2 n_3$	$e_3^t = t n_4$
$e_4^t = \frac{1}{t^4} n_4$	$e_4^t = t^2 n_4$	$e_4^t = it(n_2 - n_3)$	$e_4^t = t n_4$	$e_4^t = t n_3 + t^2 n_4$
$e_5^t = \frac{1}{t^6} n_5$	$e_5^t = t^3 n_5$	$e_5^t = t n_5$	$e_5^t = t^2 n_5$	$e_5^t = t^2 n_5$

To obtain $\mathcal{J}_{21} \to \mathcal{J}_9$ consider the following change of basis $e_1^t = -\frac{1}{\sqrt{2t}} n_1$, $e_2^t = \frac{1}{\sqrt{2t}} n_2$, $e_3^t = -\frac{1}{2t^2} n_4$, $e_4^t = -\frac{1}{\sqrt{2}} n_1 + \frac{1}{\sqrt{2}} n_2 - n_3 + \left(\frac{1}{2} - \frac{1}{4t}\right) n_4$ and $e_5^t = \frac{1}{2t^2} n_5$ of $\mathcal{J}_{21}$. Deformations $\mathcal{J}_4 \to \mathcal{J}_3 \to \mathcal{J}_2$ and $\mathcal{J}_4 \to \mathcal{J}_1$ follow from Example 6. Suppose $\{n_1, \ldots, n_5\}$ is the basis of $\mathcal{J}_9$, then $e_1^t = t n_1 - \frac{1}{2} n_3 + t^2 n_4$, $e_2^t = t n_2 - \frac{1}{2} n_3 + t^2 n_4$, $e_3^t = t^2 (n_3 - n_5)$, $e_4^t = \frac{t}{2} n_3 + t^3 n_4$ and $e_5^t = t^4 n_5$ for any $t \neq 0$ defines algebras which are isomorphic to $\mathcal{J}_9$. Meanwhile $e_3^t e_4^t = t e_5^t$ thus, when t tends to zero, we get the structure constants of algebra $\mathcal{J}_{10}$. The

basis $e_1^t = n_1 - t^2 n_2 + \sqrt{2} t n_4$, $e_2^t = -t^2 n_1 + n_3 - \sqrt{2} t n_4$, $e_3^t = t^4 n_3 - 4t^2 n_5$, $e_4^t = \frac{t^4}{2} n_3 - \frac{t^5}{2\sqrt{2}} n_4$ and $e_5^t = -t^6 n_5$ of $\mathcal{J}_{10}$ does the trick for $\mathcal{J}_{10} \to \mathcal{J}_{13}$. To complete the description of orbits inside $\mathcal{J}_{21}^{\mathrm{G}}$ use Lemma 7 for the following cases:

$\mathcal{J}_8 \to \mathcal{J}_7$	$\mathcal{J}_7 \to \mathcal{J}_4$	$\mathcal{J}_{13} \to \mathcal{J}_{14}$	$\mathcal{J}_{10} \to \mathcal{J}_5$	$\mathcal{J}_5 \to \mathcal{J}_6$
$e_1^t = n_2$	$e_1^t = n_1 + n_4$	$e_1^t = n_2$	$e_1^t = \frac{it}{\sqrt{2}}(n_1 - n_2)$	$e_1^t = n_1 + n_4$
$e_2^t = tn_1$	$e_2^t = n_2 + n_4$	$e_2^t = t(-n_1 + n_2)$	$e_2^t = t^2 n_3$	$e_2^t = n_2 + n_5$
$e_3^t = tn_3$	$e_3^t = n_3$	$e_3^t = -tn_3$	$e_3^t = \frac{t^2}{2}(n_1 + n_2)$	$e_3^t = tn_3$
$e_4^t = n_4$	$e_4^t = n_5$	$e_4^t = n_4$	$e_4^t = t^2 n_4$	$e_4^t = tn_4$
$e_5^t = tn_5$	$e_5^t = tn_1$	$e_5^t = -tn_5$	$e_5^t = t^4 n_5$	$e_5^t = tn_5$

Any algebra $\mathcal{J}_{28}$, $\mathcal{J}_{30}$-$\mathcal{J}_{34}$, $\mathcal{J}_{39}$ as well as the family $\mathcal{J}_{15}^{\alpha}$, $\alpha \in \mathbf{k}$ deforms into algebra $\mathcal{J}_{22}$. Deformations $\mathcal{J}_{22} \to \mathcal{J}_{15}^{\alpha} \to \mathcal{J}_{39}$ follow from Sections 4.1 and 4.2. To show that $\mathcal{J}_{39} \to \mathcal{J}_{34}$ we use again Lemma 7 applying it to the curve $\gamma(t) : e_1^t = n_1$, $e_2^t = tn_2$, $e_3^t = tn_3$, $e_4^t = tn_4$ and $e_5^t = n_5$ of $\mathcal{J}_{39}$. The deformation $\mathcal{J}_{22} \to \mathcal{J}_{33}$ is given by the change of basis $e_1^t = t^2 n_1 + n_3$, $e_2^t = t^4 n_2 + n_4$, $e_3^t = tn_2 - t^3 n_3$, $e_4^t = t^5 n_5$ and $e_5^t = t^6 n_4 + t^2 n_5$ of $\mathcal{J}_{22}$. The curve $\gamma(t)$:

$$
\begin{aligned}
e_1^t &= t^2 n_1 + n_2 + tn_3, \\
e_2^t &= t^4 n_2 + 2tn_4 + 3t^2 n_5, \\
e_3^t &= \tfrac{t^4}{3} n_2 + \tfrac{2t^5}{3} n_3, \\
e_4^t &= \tfrac{2t^9}{3} n_4, \\
e_5^t &= t^5 n_4 + t^6 n_5
\end{aligned}
$$

belongs to $\mathcal{J}_{33}^{\mathrm{G}}$ for any $t \neq 0$ and intersects $\mathcal{J}_{32}$ when $t = 0$, thus $\mathcal{J}_{33} \to \mathcal{J}_{32}$. The new basis $e_1^t = n_1$, $e_2^t = n_2$, $e_3^t = t(n_2 - n_3)$, $e_4^t = -tn_4$ and $e_5^t = n_5$ of $\mathcal{J}_{32}$ for any $t \neq 0$ defines algebra isomorphic to $\mathcal{J}_{32}$. Observe that $\left(e_3^t\right)^2 = 2te_4^t$, thus we get the structure $\mathcal{J}_{31}$ when t tends to zero, i.e. $\mathcal{J}_{32} \to \mathcal{J}_{31}$. The changes of basis $e_1^t = t^2 n_1 + \frac{1}{9} n_2 + \frac{t}{3} n_3$, $e_2^t = t^4 n_2 + \frac{2t}{27} n_4 + \frac{t^2}{3} n_5$, $e_3^t = \frac{1}{12} n_2 + \frac{t}{2} n_3$, $e_4^t = \frac{t^5}{2} n_4$ and $e_5^t = \frac{t}{12} n_4 + \frac{t^2}{4} n_5$ of $\mathcal{J}_{33}$ and $e_1^t = t(n_1 + n_2)$, $e_2^t = t^2 n_2$, $e_3^t = \frac{1}{2} n_2 + n_3$, $e_4^t = t^2 n_4$ and $e_5^t = n_4 + n_5$ of $\mathcal{J}_{30}$ give $\mathcal{J}_{33} \to \mathcal{J}_{30}$ and $\mathcal{J}_{30} \to \mathcal{J}_{28}$, respectively.

Next, $\mathcal{J}_{40} \to \mathcal{J}_{38} \to \mathcal{J}_{36}$. Indeed, we apply Lemma 7 to the change of basis $e_1^t = \frac{1}{t^2} n_1$, $e_2^t = \frac{1}{t} n_2$, $e_3^t = \frac{1}{t^3} n_3$, $e_4^t = \frac{1}{t^5} n_4$ and $e_5^t = \frac{1}{t^4} n_5$ of $\mathcal{J}_{40}$ and $e_1^t = n_2$, $e_2^t = tn_1$, $e_3^t = tn_3$, $e_4^t = tn_5$ and $e_5^t = t^2 n_4$, of $\mathcal{J}_{38}$, respectively.

Finally, the orbits of remaining structures, i.e. families $\mathcal{J}_{23}^{\beta}$, $\mathcal{J}_{24}^{\gamma}$, $\mathcal{J}_{26}^{\delta}$ as well as algebras $\mathcal{J}_{25}$, $\mathcal{J}_{35}$, $\mathcal{J}_{41}$, belong to the closure of $\mathcal{N}_{27}^{\#}$. The deformations $\mathcal{J}_{27}^{\varepsilon,\phi} \to \mathcal{J}_{26}^{\delta}$, $\mathcal{J}_{23}^{\beta} \to \mathcal{J}_{41}$ and $\mathcal{J}_{23}^{\beta} \to \mathcal{J}_{24}^{\gamma}$ follow from Sections 4.2 and 4.3.

Further, we show that $\mathcal{J}_{27}^{\varepsilon,\phi} \to \mathcal{J}_{23}^{\beta_0}$ for any fix $\beta_0 \in \mathbf{k}$. Consider the change of basis of $\mathcal{J}_{27}^{\varepsilon(t),\phi(t)}$:

$$
\begin{aligned}
e_1^t &= (-1 - t\beta_0) n_1 - n_2, \\
e_2^t &= tn_2, \\
e_3^t &= (1 + t\beta_0)^2 n_3 + n_4, \\
e_4^t &= -tn_4, \\
e_5^t &= t^2 (1 + t\beta_0) n_5,
\end{aligned}
\qquad (\varepsilon(t), \phi(t)) = \left(-1 - t\beta_0, \frac{-1 - t^2 - t\beta_0}{(1 + t\beta_0)^2} \right).
$$

Note that $\varepsilon(t)\phi(t) = 1 + \frac{t^2}{1 + t\beta_0}$ then, while for any $t \neq 0$ this basis defines structure in $\mathcal{N}_{27}^{\#}$, for $t = 0$ it gives $\mathcal{J}_{23}^{\beta_0}$.

Applying Lemma 7 to the basis transformation $e_1^t = n_2$, $e_2^t = tn_1 - tn_2$, $e_3^t = tn_3$, $e_4^t = tn_4$ and $e_5^t = -2t^2 n_3 + t^2 n_5$ of $\mathcal{J}_{41}$ we obtain $\mathcal{J}_{41} \to \mathcal{J}_{35}$.

To show that $\mathcal{J}_{24}^{\gamma} \to \mathcal{J}_{25}$, define a curve $\phi(t) \subset \mathcal{J}\mathrm{orN}_5$, as been the set of structure constants of algebras with the basis $e_1^t = n_1 + n_2$, $e_2^t = -2tn_2$, $e_3^t = n_3 + 2n_4$, $e_4^t = -2tn_4$ and $e_5^t = tn_5$, where $\{n_1, \dots, n_5\}$ is a basis of $\mathcal{J}_{24}^{\gamma(t)}$, $\gamma(t) = \frac{t-1}{2}$. Analogously to Section 4.2, the curve lies transversely to the orbits of $\mathcal{J}_{24}^{\gamma}$. For any $t \neq 0$, $\phi(t) \subset \mathcal{N}_{24}^{\#}$, while $\phi(0)$ corresponds to the structure constants of $\mathcal{J}_{25}$. □

Lemma 12. *The infinite union $\mathcal{N}_{27}^{\#}$ is not contained in the Zariski closure of the orbits of any algebra in $\mathcal{J}\mathrm{orN}_5$.*

Proof. By Corollary 10 the non-associative algebra $\mathcal{J}_{27}^{\varepsilon,\phi}$ can not be deformed into the associative algebra ϵ_1 for any $\varepsilon, \phi \in \mathbf{k}$. Observe that for any $\varepsilon, \phi \in \mathbf{k}$ with $\varepsilon\phi \neq 1$, $\dim \left(\mathcal{J}_{27}^{\varepsilon,\phi}\right)^2 = 3$ and $\dim \mathrm{Ann}(\mathcal{J}_{27}^{\varepsilon,\phi}) = 1$. Applying Theorem 9 item 3 and item 2 we obtain that $\mathcal{J}_{21} \nrightarrow \mathcal{J}_{27}^{\varepsilon,\phi}$ and $\mathcal{J}_{40} \nrightarrow \mathcal{J}_{27}^{\varepsilon,\phi}$, when $\varepsilon\phi \neq 1$, respectively.

To see that $\mathcal{J}_{22} \nrightarrow \mathcal{J}_{27}^{\varepsilon,\phi}$, first observe that $\mathcal{J}_{22}$ does not deform into the other three algebras from Lemma 11. Indeed, comparing the dimensions of the automorphism group we deduce that $\epsilon_1 \nrightarrow \mathcal{J}_{22}$ and $\mathcal{J}_{40} \nrightarrow \mathcal{J}_{22}$, while the dimensions of the second power provide that $\mathcal{J}_{22}^{\mathrm{G}} \not\subseteq \overline{\mathcal{J}_{21}^{\mathrm{G}}}$. Suppose that $\mathcal{N}_{27}^{\#} \subseteq \overline{\mathcal{J}_{22}^{\mathrm{G}}}$ then $\mathcal{J}_{22}$ is rigid and $\mathcal{C} = \overline{\mathcal{J}_{22}^{\mathrm{G}}}$ is a irreducible component of $\mathcal{J}\mathrm{orN}_5$. Since $\overline{\mathcal{N}_{27}^{\#}}$ is contained in the irreducible component $\mathcal{C}$ and both has the same dimension $\dim \overline{\mathcal{N}_{27}^{\#}} = 21 = \dim \mathcal{C}$, $\overline{\mathcal{N}_{27}^{\#}} = \mathcal{C}$. So, $\mathcal{N}_{27}^{\#}$ is a dense subset of $\mathcal{C}$ and $\mathcal{J}_{22}^{\mathrm{G}}$ is open in $\mathcal{C}$ then there exists $\mathcal{J} \in \mathcal{N}_{27}^{\#}$ such as $\mathcal{J} \in \mathcal{J}_{22}^{\mathrm{G}}$, i.e., $\mathcal{J} \simeq \mathcal{J}_{22}$, a contradiction by Theorem 4.

□

Corollary 13. *$\overline{\mathcal{N}_{27}^{\#}}$ determines a component of the variety $\mathcal{J}\mathrm{orN}_5$. In particular, we proved that a finite-dimensional Jordan algebras not necessarily deforms into rigid one.*

Proof. From Lemma 12 it follows that we only have to show that all algebras in $\overline{\mathcal{N}_{27}^{\#}}$ belong to the same component. In fact $\overline{\mathcal{N}_{27}^{\#}}$ is a product of two irreducible varieties $\left(\mathcal{J}_{27}^{\varepsilon,\phi}\right)^{\mathrm{G}}$ and $\mathbf{k}^2 = \{(\varepsilon, \phi) \,|\, \varepsilon, \phi \in \mathbf{k}\}$ and is therefore irreducible. □

Lemma 14. *Algebras ϵ_1, $\mathcal{J}_{21}$, $\mathcal{J}_{22}$ and $\mathcal{J}_{40}$ are rigid.*

Proof. By a similar argument as in proof of Lemma 12 we can show that $\mathcal{J}_{22}^{\mathrm{G}} \not\subseteq \overline{\mathcal{N}_{27}^{\#}}$ and therefore $\mathcal{J}_{22}$ is rigid. To show that the other three algebras are rigid it is sufficient to prove that there are no deformations between them and that none of these algebras is dominated by the family $\mathcal{J}_{27}^{\varepsilon,\phi}$.

Using the dimension argument for the second power of an algebra we have $\mathcal{J}_{21} \nrightarrow \epsilon_1$, $\mathcal{J}_{22} \nrightarrow \epsilon_1$ and $\mathcal{J}_{21} \nrightarrow \mathcal{J}_{40}$, while comparing the dimension of the annihilator we obtain $\mathcal{J}_{40} \nrightarrow \epsilon_1$. Thus we have shown that ϵ_1 is a rigid algebra in $\mathcal{J}\mathrm{orN}_5$ too. The associative algebra ϵ_1 can not dominate non-associative algebras, thus $\epsilon_1 \nrightarrow \mathcal{J}_{21}$ and $\epsilon_1 \nrightarrow \mathcal{J}_{40}$. Also, due to the dimension argument for the automorphism group we get $\mathcal{J}_{22} \nrightarrow \mathcal{J}_{40}$ and comparing the dimension of the associative center we have $\mathcal{J}_{22} \nrightarrow \mathcal{J}_{21}$ and $\mathcal{J}_{40} \nrightarrow \mathcal{J}_{21}$. On the other hand $\dim \mathcal{J}_{21}^{\mathrm{G}} = 22$ while $\dim \overline{\mathcal{N}_{27}^{\#}} = 21$ leading to $\mathcal{J}_{21}^{\mathrm{G}} \not\subseteq \overline{\mathcal{N}_{27}^{\#}}$ and therefore $\mathcal{J}_{21}$ is a rigid algebra. Finally suppose that $\mathcal{J}_{40}$ belongs to the component $\overline{\mathcal{N}_{27}^{\#}}$. Note that the dimensions of $\mathcal{J}_{40}^{\mathrm{G}}$ and $\mathcal{N}_{27}^{\#}$ are both equal to 21, therefore

$$\overline{\mathcal{J}_{40}^{\mathrm{G}}} = \overline{\mathcal{N}_{27}^{\#}}.$$

Which is impossible since $\dim \mathrm{Ann}(\mathcal{J}_{40}) = 2$, while $\dim \mathrm{Ann}(\mathcal{J}_{27}^{\varepsilon,\phi}) = 1$ for $\varepsilon, \phi \in \mathbf{k}$ with $\varepsilon\phi \neq 1$. Consequently $\mathcal{J}_{40}$ is a rigid algebra. □

Finally, note that the change of basis given by identity matrix I_5 multiplied by t guarantees $\mathcal{J} \to \epsilon_{25}$ for any $\mathcal{J} \in \mathcal{J}\mathrm{orN}_5$. Consequently, $\mathcal{J}\mathrm{orN}_5$ is a connected affine variety. Therefore we have proved the following theorem.

Theorem 15. *The algebraic variety $\mathcal{J}\mathrm{orN}_5$ of five-dimensional nilpotent Jordan algebras is a connected affine variety with an infinite number of orbits under the action of the group G. Furthermore, it has four irreducible components given by Zariski closures of the orbits of the rigid algebras ϵ_1, $\mathcal{J}_{21}$, $\mathcal{J}_{22}$ and $\mathcal{J}_{40}$ and one more component that is the Zariski closure of the infinite union $\mathcal{N}_{27}^{\#}$.*

The diagram of $\mathcal{J}\mathrm{orN}_5$ was represented in Figure 1.



Figure 1: The irreducible components of $\mathcal{J}\mathrm{orN}_5$

6 References

[1] F. Flanigan, Algebraic Geography: Varieties of Structure Constants, Pacific Journal of Mathematics 27 (1) (1968) 71–79.

[2] G. Mazzola, Generic Finite Schemes and Hochschild Cocycles, Commentarii Mathematici Helvetici 55 (1980) 267–293.

[3] A. Makhlouf, The irreducible components of the nilpotent associative algebras, Revista Matemática Complutense 6 (1) (1993) 27.

[4] F. Grunewald, J. O'Halloran, Varieties of nilpotent Lie algebras of dimension less than six, Journal of Algebra 112 (2) (1988) 315–325.

[5] S. Craig, Degenerations of 6-dimensional nilpotent lie algebras over $\mathbb{C}$, Communications in Algebra 18 (10) (1990) 3493–3505.

[6] M. Goze, J. A. Bermudez, On the varieties of nilpotent lie algebras of dimension 7 and 8, Journal of pure and applied algebra 77 (2) (1992) 131–140.

[7] J. M. Ancochea-Bermudez, J. R. Gómez-Martin, G. Valeiras, M. Goze, Sur les composantes irréductibles de la varieté des lois d'algèbres de lie nilpotentes, Journal of Pure and Applied Algebra 106 (1) (1996) 11 – 22.

[8] I. Kashuba, I. Shestakov, Jordan Algebras of Dimension Three: Geometric Classification and Representation Type, Actas del XVI Coloquio Latinoamericano de Álgebra (Colonia del Sacramento, Uruguay). Biblioteca de la Revista Matemática Iberoamericana 1 (2005) 295–315.

[9] I. Kashuba, M. E. Martin, Deformations of Jordan Algebras of Dimension Four, Journal of Algebra 399 (2014) 277–289.

[10] J. M. Ancochea Bermúdez, J. Fresán, J. M. Bentabol, Contractions of Low-Dimensional Nilpotent Jordan Algebras, Communications in Algebra 39 (3) (2011) 1139–1151.

[11] M. E. Martin, Four Dimensional Jordan Algebras, International Journal of Mathematics, Game Theory and Algebra 20 (4) (2013) 41–59.

[12] A. Hegazi, H. Abdelwahab, Classification of five dimensional nilpotent jordan algebras, arXiv preprint arXiv:1401.3992.